

EDITORIAL REVIEW

Thromboxane synthesis and action within the kidney

Thromboxane (Tx) A₂ is a labile derivative of arachidonic acid metabolism that amplifies the platelet response to a variety of aggregating agents [1]. In addition, TxA₂ contracts vascular smooth muscle [2] and glomerular mesangial cells [3] through receptor-mediated actions, resulting in increased intracellular calcium [4–6]. TxA₂ is formed by the sequential cyclooxygenation and peroxidation of arachidonic acid to the prostaglandin (PG) endoperoxides, PGG₂ and PGH₂. The last of these compounds is then transformed to TxA₂ by a specific TxA₂ synthase.

Platelets represent a major source of TxA₂ formed *in vivo* under physiological conditions. However, TxA₂ synthase is widely distributed in other cells, particularly in macrophages, in the lung, peritoneum and kidney [7, 8], which may augment their contribution to systemic TxA₂ biosynthesis in disease states. The most convincing evidence for the biological importance of TxA₂ and its precursor endoperoxides derives from the beneficial effects of aspirin in platelet-mediated vascular occlusive disease [9]. Aspirin irreversibly inhibits platelet cyclooxygenase activity by acetylating a serine residue of the enzyme PGG/H synthase [10, 11]. This mechanism is presumed to underlie its effectiveness in preventing death in unstable angina [12–14] and in patients who have suffered a myocardial infarction [15], conditions in which TxA₂ biosynthesis is augmented [16].

More recently, evidence has accumulated to support a role for these eicosanoids in the pathogenesis of various forms of renal disease. The purpose of this review will be to consider the role of TxA₂ within the kidney and to consider the potential therapeutic benefit which might derive from modulating TxA₂ synthesis and/or action by pharmacological or dietary intervention.

Mechanism of action of TxA₂

Due to the evanescence of TxA₂, which has a half-life of 30 seconds at physiological pH [1, 17], the receptor site for this compound has been characterized using structural analogs of PGH₂ and, more recently, TxA₂ itself [18–20]. Indeed, the chemical structure of TxA₂ was at first merely hypothesized and only confirmed 10 years later when a TxA₂ macrolactone was synthesized and the effects on platelets and vascular smooth muscle of the biologically generated material were replicated [17]. Given the limitation of the ligands used to characterize PGH₂/TxA₂ receptor sites, it has been thought that both eicosanoids have a similar affinity for a common receptor site on platelets [21].

Recent observations suggest that the situation might be more complex. Thus, the antagonist ligand, GR 32191 [22], identifies two sites on human platelets; it dissociates rapidly from one (GR_r) and appears to bind irreversibly (GR_{irr}) to the other [23]. GR_{irr} sites are coupled to phospholipase C and transduce diacylglycerol formation, phosphoinositide turnover, an increase of intracellular calcium and activation of protein kinase C. These events are associated with platelet activation and secretion. GR_r sites, by contrast, mediate PGH₂/TxA₂-evoked platelet shape change and a major portion of the ligand-induced increase in intracellular calcium, presumably from an IP₃-insensitive source [23]. Discrimination of the shape change and aggregation responses was also achieved with the reversibly bound agonist ligand, BOP [24], although this observation in isolation might have been explained by varied degrees of receptor occupancy necessary to evoke the aggregation and shape change responses [25]. However, this seems an unlikely explanation, given the marked distinction between the plasma half-life of GR 32191 (6 to 7 hrs) and the time to recovery of the ligand-evoked aggregation response *ex vivo*, which corresponds to platelet turnover time (8 to 10 days). Thus, the pattern of inactivation of PGH₂/TxA₂ receptors on human platelets is consistent with the existence of distinct subtypes of the receptor. It is unknown whether these sites represent two distinct receptor isoforms or varied degrees of posttranslational modification of a single receptor or if the two sites discriminated by GR 32191 differ in their affinities for PGH₂ and TxA₂ [26].

The sequence of a full length cDNA for a PGH₂/TxA₂ receptor obtained from human placenta has recently been reported [27]. Hydrophobicity plotting of the deduced amino acid sequence suggests that the PGH₂/TxA₂ receptor consists of seven transmembrane spanning domains, characteristic of the G protein-linked receptor superfamily. The identity between this clone and a partial length cDNA from a human megakaryocytic cell line was used to suggest the identity of vascular (that is, placental) and platelet (that is, megakaryocyte-derived) PGH₂/TxA₂ receptors [27]. Further information at the molecular level will be necessary to determine if this hypothesis, which is contrary to much biochemical evidence and precedent with most other receptors, is supported.

Studies of the respective hierarchies for structurally distinct agonist and antagonist ligands in platelets and vascular smooth muscle cells suggest that tissue specific heterogeneity is likely [28–30]. However, others who have performed similar experiments, have disputed this interpretation of their data [31–33]. Recently, additional evidence that supports the existence of tissue specific heterogeneity of PGH₂/TxA₂ receptors has been provided [34]. Thus, dissociation of GR 32191 from vascular smooth muscle cells, derived from both rat aorta and human saphenous vein, identifies only a single site, from which the ligand dissociates rapidly. In addition, we have noted that platelet activating factor (PAF) heterologously down regulates

Received for publication January 18, 1991

and in revised form April 8, 1991

Accepted for publication April 12, 1991

© 1992 by the International Society of Nephrology

GR_{irr} but not GR_r sites on human platelets [35]. Consistent with the absence from vascular smooth muscle cells of the subtype that mediates aggregation in platelets, agonist evoked responses in this tissue are not down regulated by PAF [34]. Radiolabeled ligands have identified a single class of high-affinity specific binding sites in glomeruli, but no studies have been reported to date with GR 32191.

PGH₂/TxA₂ analogs activate phospholipase C in mesangial cells in a manner analogous to that observed in platelets. They cause an immediate increment in calcium derived from intracellular stores followed by a sustained elevation for longer than 10 minutes, which is dependent upon extracellular calcium. These effects are blocked by a PGH₂/TxA₂ antagonist. Studying this action of the PGH₂/TxA₂ mimetic, U46619 and PGI₂ analog, ilprost, on proliferation of cultured human mesangial cells in the presence or absence of growth factors, Mene et al found multiple functional interactions between eicosanoids and growth factors in the control of mesangial cell proliferation [36]. In particular, U46619 stimulates proliferation when used alone but inhibits proliferation when applied concomitantly with growth factors.

The biochemical mechanisms which regulate the response to PGH₂/TxA₂ have been most clearly defined in platelets. Homologous desensitization of the aggregation response to PGH₂/TxA₂ agonists occurs rapidly, with a half-time of two to three minutes. This results in uncoupling of the aggregation receptor from its phospholipase C-linked G protein [4], with subsequent loss of receptors from the cell membrane. The nature of the G protein involved is unknown, although it is insensitive to pertussis toxin [37], in contrast to that by which thrombin and PAF activate phospholipase C [37–40]. Coincident with these events, the platelet adenylate cyclase is sensitized to stimulation at the post-receptor level [35]. This effect is transduced through the GR_{irr} receptor. PGH₂/TxA₂ released from activated platelets stimulates PGI₂ generation by adjacent endothelial cells [41]. Cross talk between the phospholipase C and adenylate cyclase pathways within thromboxane stimulated platelets would serve to amplify the inhibitory effects of PGI₂. Finally, despite the transduction of these effects via the GR_{irr} receptor and some tenuous evidence that the PGH₂/TxA₂ receptor is phosphorylated upon desensitization [42], neither desensitization of aggregation, nor sensitization of the cyclase, appear dependent on protein kinase C [35].

PGH₂/TxA₂ evoked intracellular calcium responses in human mesangial cells through activation of phospholipase C also undergo rapid, homologous desensitization [6]. In this system, the biochemical mechanisms which underlie this effect and the possible cross talk with other systems remain to be defined. The density of PGH₂/TxA₂ binding sites is reduced in mesangial cells obtained from rats made diabetic with streptozotocin [43], a condition associated with enhanced glomerular TxA₂ synthesis [44].

Evidence for TxA₂ synthesis by resident renal cells

The hypothesis that resident renal cells can metabolize arachidonic acid to prostaglandins and TxA₂ was based on the observation that glomerular capillaries have immunoreactive cyclooxygenase activity [45]. Soon after this observation, two independent laboratories have documented that isolated rat glomeruli convert arachidonic acid into prostaglandins and

TxA₂ [46, 47]. Subsequent studies documented that eicosanoid biosynthesis occurs in glomeruli of other species, including humans [48–50]. Similar conclusions were derived from studies on renal epithelial and mesangial cells in culture [51, 52]. PGH₂/TxA₂ evoked intracellular calcium responses in human mesangial cells through activation of phospholipase C also undergo rapid, homologous desensitization [6].

Angiotensin II and arginine-vasopressin stimulate glomerular TxA₂ synthesis via specific receptor(s) linked to phospholipase C [53]. More recently, endothelin has been found to induce the synthesis of TxA₂ by rat and bovine mesangial cells [54, 55]. Indeed studies with antagonists suggest that PGH₂/TxA₂ receptors mediate the contractile response to endothelin in guinea pig tracheal strips and rat aorta [56, 57]. In some [54], but not all [55] systems, the hormonal stimulation of mesangial cell TxA₂ can be blocked by lowering the concentration of extracellular calcium or by incubating the cells with the calcium antagonists, verapamil or nifedipine.

Factors other than hormones which regulate renal TxA₂ biosynthesis include cell-derived mediators, like platelet activating factor (PAF) and interleukin 1 (IL-1), complement membrane attack complex, immune complexes and endotoxin. PAF induces TxA₂ synthesis by rat mesangial cells in culture through a receptor-mediated effect [58]. IL-1 stimulates TxA₂ release after continuous, prolonged exposure to rat mesangial cells [59]. This is suggestive of an effect dependent upon new protein synthesis, perhaps due to regulation of cyclooxygenase message, as has been shown in dermal fibroblasts and endothelial cells [60, 61]. Oxygen-derived free radicals also activate phospholipase A₂, thus enhancing prostaglandin and TxA₂ synthesis in isolated glomeruli through an increase in the availability of arachidonic acid [62]. Cultured mesangial cells exhibit a biphasic response to progressively higher concentrations of hydrogen peroxide. The decrease in eicosanoid formation at higher concentrations is probably due to inactivation of cyclooxygenase [63]. An increase in TxA₂ production occurs in rat glomerular epithelial cells when the terminal complement components C5b-9 are added to the culture medium [64]. *E. coli* lipopolysaccharide stimulates both PGE₂ and TxA₂ synthesis by cultured mesangial cells in a dose- and time-dependent manner [65]. Exposure of mesangial cells to immune complexes [66] or the binding of an antibody to a cell membrane antigen [67] also promotes arachidonic release and eicosanoid formation in mesangial cells.

Another important pathway which may result in renal generation of TxA₂ has been reported by Sraer et al [68] who demonstrated that human glomeruli released a diffusible factor capable of stimulating platelet TxA₂ synthesis. The clinical nature of this factor has not yet been fully identified. GC/MS analysis suggests that at least part of the stimulatory activity is attributable to saturated and monoenoic acids. Thus, injury to glomerular cells may activate platelets by a pathway which is independent from the formation of collagen and leads to local formation of TxA₂.

Modulation of renal function by TxA₂ in health and disease

Whole-kidney clearance studies in rats and dogs have shown a consistent fall in glomerular filtration rate (GFR) and in renal plasma flow (RPF) in response to systemic infusion of stable prostaglandin endoperoxide analogues [69–71]. Micropuncture

A Control



B Infusion of the TxA_2 mimetic U-46619

Fig. 1. Schematic representation of changes in single nephron glomerular filtration rate and its determinants during infusion of the TxA_2 mimetic U-46619 in euvoletic rats. (a), from reference 76 and (b) from reference 69. Abbreviations are: ΔP , mean glomerular transcapillary hydraulic pressure gradient; K_f , glomerular capillary ultrafiltration coefficient; Q_a , glomerular capillary plasma flow rate; R_a and R_e , afferent and efferent arteriolar resistances; SNGFR, single nephron glomerular filtration rate.

studies performed in rats during intraarterial infusion of the $\text{PGH}_2/\text{TxA}_2$ analog, U46619, have shown a rise in afferent and efferent arteriolar resistance with a parallel reduction of glomerular capillary flow and single nephron GFR [69] (Fig. 1). Moreover, in an isolated perfused rat kidney preparation, the analog decreased both RPF and GFR. However, the decrease in GFR was more marked than in RPF, suggesting that the mimetic may act selectively upon afferent resistance vessels or may reduce GFR by lowering the ultrafiltration coefficient (K_f) [72].

More recently, this possibility has been addressed directly by examining the effect of $\text{PGH}_2/\text{TxA}_2$ analogs on renal microvessels. This has been done in a hydronephrotic perfused kidney model that allows direct visualization of the renal microcirculation [73]. Image analysis of afferent and efferent arteriolar diameters provided direct evidence that $\text{PGH}_2/\text{TxA}_2$ preferentially constricts the afferent arteriole. Of interest, is the observation that such a constrictor effect of thromboxane was abolished by calcium antagonists [72]. No changes in K_f were detected since the single nephron GFR and glomerular capillary plasma flow rate (Q_a) fell in parallel. However, most rats in this study were evaluated at filtration pressure equilibrium during infusion of the $\text{PGH}_2/\text{TxA}_2$ analog, so that it was not possible to calculate exact values of K_f . Thus, small TxA_2 -mediated changes in K_f may have gone undetected.

Consistent with this possibility, GFR falls by a greater extent than RPF in the early phase of nephrotoxic nephritis characterized by enhanced glomerular TxA_2 synthesis, suggesting a fall in K_f [74]. Similarly, direct measurements of glomerular hemodynamic changes during infusion of antiglomerular basement membrane antibody [75] indicated a fall in K_f together with an increase in intraglomerular capillary pressure. On the other hand, a direct effect of TxA_2 on K_f is consistent with *in vitro* data showing that endoperoxide analogs contract rat isolated glomeruli as well as cultured mesangial cells, thereby presumably reducing the filtration surface area *in vivo* [6, 76].

Recent data have indicated an interesting strain difference

between young SHR and age matched WKY in the renal vascular reactivity to locally administered angiotensin II and TxA_2 [77]. Unlike WKY, SHR display an exaggerated vascular reactivity to both angiotensin II and TxA_2 , indicating that the degree of renal vasoconstriction in response to agonists may vary across strains of animals in a way that is genetically determined. Whether this would contribute to the sustained renal vasoconstriction of animals which are hypertensive remain to be established.

Angiotensin II, which induces TxA_2 synthesis by mesangial cells, also induces a concentration dependent increase in PGE_2 [52] and PGI_2 synthesis [78] that causes mesangial relaxation by promoting an increase in cyclic AMP. This modulating effect of PGE_2 and PGI_2 is lost in the presence of cyclooxygenase inhibitors, so that the mesangial response is amplified and GFR decreases. Similarly, whole glomeruli decrease their surface area when incubated with angiotensin II, and this effect is exacerbated by prostaglandin inhibition [76].

In a rat model of glycerol-induced acute renal failure the glomerular synthesis of PGE_2 decreased in the early phase of the disease while TxA_2 synthesis was greater in glycerol treated than in control rats only upon stimulation with arachidonic acid and at 24 hours after the glycerol administration [79]. These findings were consistent with other data showing that an inhibitor of TxA_2 synthesis partially protected another strain of rats against glycerol-induced acute renal failure [80]. Of interest, an angiotensin converting enzyme inhibitor did not confer protection in the same setting, but rather resulted in a significant increase in renal TxB_2 and 6-keto- $\text{PGF}_{1\alpha}$ synthesis. This may indicate that, at least in this model, TxA_2 is an important mediator of damage.

In a more physiological setting, the administration of cyclooxygenase inhibitors to conscious, sodium-replete animals does not induce changes in RPF and GFR, indicating that a compensatory role of vasodilatory prostaglandins for maintaining renal function is negligible under these circumstances [81–83]. By contrast, in anesthetized [84] or sodium-deprived animals [81],

or those with hemorrhagic hypotension [85] or congestive heart failure [86] blocking the cyclooxygenase leads to a fall in RPF and GFR. Thus, the potentially adverse effects of increased production of mesangial cell contractile agonists, including $\text{PGH}_2/\text{TxA}_2$, is partially offset by relaxant eicosanoids such as PGE_2 and PGI_2 . Neither TxA_2 , nor the other eicosanoids, appear to play an important role in mesangial function under physiological circumstances. Thus, it is not surprising that a thromboxane synthase inhibitor does not alter p-aminohippurate or inulin clearance in healthy humans [87], while a thromboxane receptor antagonist is capable of increasing RPF and GFR in patients with lupus nephritis [88].

Measurement of TxA_2 biosynthesis

Because of chemical instability of the bicyclic oxane structure, TxA_2 is rapidly hydrolyzed to the chemically stable and biologically inactive derivative, TxB_2 . Thus, in vivo as well as in vitro measurements of TxA_2 biosynthesis have relied upon detection of TxB_2 or its enzymatic metabolites [89]. In healthy human beings, TxB_2 is produced at a very low rate, that is, $0.11 \text{ ng/kg min}^{-1}$ [90], possibly reflecting the low frequency and/or intensity of stimuli to its secretion under physiologic circumstances. In contrast to the low rate of actual in vivo production, the capacity of platelets to synthesize TxA_2 ex vivo in response to appropriate stimuli (for example, thrombin) is quite substantial; 1 ml of human whole blood incubated at 37°C for one hour producing almost as much TxB_2 as the whole body during the same period of time, that is, roughly 300 to 400 ng [90]. TxB_2 undergoes two major pathways of metabolism in humans [91], one involving β -oxidation resulting in the formation of 2,3-dinor- TxB_2 and the other involving dehydrogenation of the hemiacetal alcohol group at C-11, resulting in the formation of a series of metabolites with a delta-lactone ring structure (such as 11-dehydro- TxB_2). These enzymatic changes are thought to occur primarily in the liver, lungs and kidney. A variety of techniques have been developed to quantitate the plasma levels and/or the urinary excretion of these metabolites [92, 93]. These include radioimmunoassay (RIA), enzyme-immunoassay (EIA) and negative ion-chemical ionization gas chromatography/mass spectrometry (NICI-GC/MS).

Largely indirect evidence supports the notion that the urinary excretion of primary unmetabolized eicosanoids, including TxB_2 , reflects intrarenal eicosanoid production under physiological circumstances [89] in contrast to the metabolites which reflect mainly systemic, extrarenal production [89, 94]. The evidence concerning urinary TxB_2 derives from infusion studies of exogenous TxB_2 in healthy subjects [90], from the investigation of patients with lupus nephritis [88] and from pharmacologic studies of selective and nonselective cyclooxygenase inhibitors in health and disease [88, 95–97]. These human studies have measured the urinary excretion of TxB_2 , and 2,3-dinor- TxB_2 and have demonstrated a dissociation of their excretory pattern (Table 1) in the setting of renal disease. No information of a similar nature is available from animal studies, even though measurements of urinary TxB_2 have been used extensively in rat and murine models of renal disease, as detailed below. Our recent finding that the rat kidney has the capacity to metabolize TxB_2 to 2,3-dinor- TxB_2 and that both are excreted in the urine, implies that under conditions of enhanced intrarenal synthesis of TxA_2 , one should expect

Table 1. Differential pattern of excretion of thromboxane B_2 and 2,3-dinor-thromboxane B_2 in health and disease

Variable	Urinary excretion	
	Thromboxane B_2	2,3-Dinor-thromboxane B_2
Thromboxane B_2 infusion at $0.1 \text{ ng/kg} \cdot \text{min}^{-1}$	no change	time-dependent increase
Aspirin 20–160 mg/day	no change	dose-dependent decrease
Sulindac 200–800 mg/day	no change	dose-dependent decrease
Lupus nephritis	2- to 5-fold increase	no change

Based on data from references 89–91 and 95–97.

increased urinary excretion of both TxB_2 and 2,3-dinor- TxB_2 [98]. Correspondingly, while urinary TxB_2 may largely derive from the kidney under physiological circumstances, augmented extrarenal TxA_2 biosynthesis would be expected to be reflected by increased excretion of TxB_2 as well as by its metabolites. Thus, urinary TxB_2 (and its metabolites) is increased in patients with platelet activation [99] and following both inhalation and ingestion of TxB_2 [100]. Whether proteinuria alters the disposition of glomerular TxA_2 or its availability for further hydrolysis in urine remains to be determined.

Measurement of ex vivo glomerular TxB_2 production in response to added substrate or other stimuli represents a capacity-related index, conceptually similar to the measurement of TxB_2 production during whole blood clotting [101]. Measurement of TxB_2 synthesis in isolated glomeruli has been used extensively to monitor disease-associated changes in $\text{PGH}_2/\text{TxA}_2$ metabolism. However, the relationship of enhanced glomerular TxB_2 production, as detected ex vivo, to the actual rate of TxA_2 biosynthesis in vivo remains to be defined. Measurement of eicosanoid production in isolated glomeruli may provide useful information on qualitative differences in disease-related expression of PG-endoperoxide metabolism, possibly reflecting changes in the cellular source(s) of TxA_2 production or upregulation of Tx-synthase gene expression.

Inhibition of TxA_2 synthesis and action

Effects on glomerular permeability to macromolecules

Little information is available thus far as to whether the course of renal disease may be modified by preventing TxA_2 synthesis or action. However, several recent studies in animal models and limited clinical investigation suggests that indices of disease severity may be modified by this type of intervention.

Some of the most encouraging information has derived from studies of drug-induced nephrosis. Thus, treatment with a selective TxA_2 synthase inhibitor was shown to reduce urinary protein excretion without changes in GFR in rats with adriamycin (ADR) induced nephrosis [102]. Whether this reflected a direct effect of TxA_2 on glomerular permeability to proteins or was the result of changes in glomerular hemodynamics induced by exaggerated renal TxA_2 synthesis was not established by that study. That pharmacological inhibition of thromboxane synthase ameliorates proteinuria in experimental nephrosis has

been confirmed [103, 104] using structurally distinct thromboxane synthase inhibitors in a different model of drug-induced (aminonucleoside) nephrosis. Thromboxane synthase inhibitors also reduced [105] membrane attack complex (MAC)-induced proteinuria in the isolated perfused model of passive Heymann nephritis. In this model, proteinuria develops acutely, making it unlikely that compositional changes in the basement membrane underlie the observation. Rather, MAC-mediated cytotoxicity of epithelial cells may explain proteinuria. Similar to the results reported earlier in ADR nephrosis, a thromboxane-synthesis inhibitor diminished proteinuria in both cases; however, the effect of thromboxane synthesis inhibition on proteinuria was only partial. Whether this is due to the fact that these compounds do not fully suppress the synthesis of thromboxane or that factors other than TxA_2 may be involved in mediating the altered glomerular permeability to proteins is still a matter of speculation. At least one study in humans has documented that inhibiting thromboxane synthesis may reduce proteinuria and induce remission of the nephrotic syndrome when it is resistant to conventional therapy [106].

A recent study of rats made diabetic with streptozotocin also showed that a thromboxane synthase inhibitor given for six months reduced urinary TxB_2 excretion by 80% and abolished the rise in urinary albumin that occurs with time in diabetes [107]. The thromboxane inhibitor also induced an increase in the clearance of inulin in diabetic animals as compared with controls. A similar study has shown that the exacerbation of proteinuria in streptozotocin rats by a high protein diet is ameliorated by thromboxane synthase inhibition [108]. No changes in blood pressure or inulin clearance were detected, suggesting a mechanism unrelated to the vasoconstrictive properties of TxA_2 .

The relationship between enhanced glomerular thromboxane formation and the development of renal disease has also been recently explored in normotensive rats of the Milan Strain (MNS) that spontaneously develop an age-related form of proteinuria and progressive glomerulosclerosis [109]. When these animals were given a thromboxane synthase inhibitor, decreased glomerular TxB_2 production and proteinuria were observed together with substantial preservation of RPF and GFR [110]. Blood pressure was unaltered by the treatment; however, the number of sclerotic glomeruli was reduced.

In nephritis induced by anti-GBM antibody, inhibitors of thromboxane synthase reduced proteinuria and partially prevented deterioration of renal function in both rat [111] and rabbit models [112]. Similar results were obtained by the use of a PAF receptor antagonist. Since the PAF receptor antagonist depressed the increment in thromboxane synthesis in immunologically injured glomeruli, it was suggested that its beneficial effect is mediated by suppression of excessive intrarenal synthesis of TxA_2 [112].

These biochemical and pharmacological findings appear to establish that exaggerated renal thromboxane synthesis may impair the selective properties of glomerular capillaries. However, the mechanism(s) for such an effect are less clear. TxA_2 may increase glomerular permeability to proteins by a direct, albeit still ill-defined, effect on glomerular barrier size-selective properties. Indirect evidence in support of this hypothesis is the observation that in ADR nephrotic rats, the increased glomerular production of TxA_2 [102] is associated with alteration of

both size and charge-dependent permeability of the glomerular barrier, as indicated by the enhanced fractional clearance of both neutral and sulfate dextrans as compared to control animals [113, 114]. Since micropuncture studies in these animals have shown no changes in the determinants of single nephron GFR [115, 116], it is tempting to speculate that TxA_2 can directly alter the selective properties of the glomerular capillary barrier. This possibility is further supported by the finding that part of the effect of PAF on glomerular membrane permeability is mediated via the secondary release of TxA_2 . Another possibility is that TxA_2 functions as an intracellular messenger for the induction of epithelial cell morphological changes that are associated with leakage of protein through the glomerular capillary wall.

Effects on K_f

In nephrotoxic serum nephritis induced by anti-GBM antibodies, glomerular TxB_2 synthesis is enhanced coincident with the reduction in GFR [111]. Blocking thromboxane synthesis by two different inhibitors induced a partial recovery of GFR, suggesting an effect of TxA_2 on factors that regulate glomerular filtration [111]. Since GFR decreased more than RPF, it is likely that the K_f is impaired by anti-GBM serum. The theoretical possibility that blocking thromboxane synthesis is beneficial in anti-GBM, because it prevents a TxA_2 -mediated decrease in K_f , is consistent with the potent contractile effect of $\text{PGH}_2/\text{TxA}_2$ mimetics on mesangial cells. This is likely to decrease K_f by reducing the filtration surface area [117].

Another example that indicates a possible role of TxA_2 in decreasing the glomerular filtration surface area in renal disease is provided by the acute toxicity of cyclosporin A (CsA). Animals and humans given CsA show a reduction in GFR associated with a progressive rise in the renal synthesis of TxA_2 [118–121]. The administration of a TxA_2 synthesis inhibitor results in a significant decrease in the urinary excretion of TxB_2 and an increase in GFR in the rat [119]. Enhanced renal synthesis of TxA_2 in CsA-treated animals correlates with the reduction in GFR but not RPF. This is consistent with the possibility that TxA_2 reduces the K_f by reducing glomerular surface area.

Effects on afferent and efferent arterioles

Besides having an effect on K_f , possibly mediated by a contractile action on mesangial cells, there is evidence that excessive renal TxA_2 synthesis can influence the tone of afferent and efferent arterioles in renal disease. In a rat model of endotoxin-induced acute renal failure, the acute increase in renal vascular resistance and decrease in RPF, GFR and filtration fraction were prevented by a thromboxane synthesis inhibitor which also abolished the excessive renal synthesis of TxA_2 [122]. The simultaneous decrease in RPF and GFR and the effect of the inhibitor in such a model are more consistent with a TxA_2 -mediated effect on afferent and efferent arterioles than on K_f .

Increased renal production of TxA_2 appears also to mediate the glomerular vasoconstriction that is associated with persistent ureteral obstruction [123]. Treatment with a thromboxane synthesis inhibitor reverses the vasoconstriction and improves renal function in this model [124]. Inflammatory cells generate excessive amounts of peptidoleukotrienes (LTs) in ex vivo

perfused hydronephrotic kidneys and both endogenous and exogenous LTC₄ stimulates TxA₂ synthesis in the hydronephrotic but not in the contralateral kidney [125]. Renal vasoconstriction is abolished by pretreatment with either a thromboxane synthase inhibitor or a leukotrene receptor antagonist [125]. Evidence that excessive renal TxA₂ synthesis mediates impaired renal function has also been obtained in autoimmune lupus mice by showing increased intrarenal production of TxB₂ in parallel deterioration of renal function, as well as by dietary manipulation of this pathway [126, 127]. Moreover, Spurney et al [127] have found that TxA₂ receptor blockade increases GFR in MRL-lpr mice.

Impairment of renal function in human lupus nephritis may also be reversed by thromboxane antagonists. Pierucci et al [88] reported that a TxA₂ receptor antagonist but not low-dose aspirin induced a similar increase in GFR and RPF in such patients, consistent with an effect of enhanced TxA₂ either on afferent and efferent arterioles or on mesangial cells. It has also been documented that the intense vasoconstriction which is associated with rejection of renal allografts is associated with increased renal TxA₂ synthesis [128]. Treatment with a thromboxane synthase inhibitor was associated with a decrease in urinary TxB₂ and an increase in RPF and GFR. Since cyclophosphamide also decreased urinary TxB₂, a likely source of the increment in TxB₂ in this model is from infiltrating inflammatory cells.

Effects on progression of glomerular damage

In the last ten years, several studies have addressed the progressive nature of renal disease after renal mass has been reduced surgically. The most attractive hypothesis is that the increased glomerular pressure and flow that develops in response to the initial loss of functioning nephrons causes glomerular injury and that this eventually leads to glomerulosclerosis [129, 130]. Rats with reduced renal mass also have increased urinary excretion of TxB₂ and chronic administration of a thromboxane synthase inhibitor decreases urinary protein excretion and limits glomerular damage [131]. The thromboxane inhibitor in this model also markedly inhibited 'in vitro' platelet aggregation, suggesting that intrarenal platelet activation and glomerular thrombosis may play a role in the progression of the disease after a critical number of nephrons has been reduced. However, the TxA₂ synthesis inhibitor also lowered blood pressure, leaving open the possibility of an effect related to this property of the compound. Indeed, neither low dose aspirin nor a thromboxane receptor antagonist—both of which failed to influence hemodynamics—retarded progressive renal disease in the remnant kidney model [132, 133].

In another model of progressive renal insufficiency that shares many biologic similarities with the human condition, immune complex nephritis induced by ferritin-anti-ferritin complexes in Dahl-salt sensitive rats, a thromboxane synthase inhibitor, ameliorated urinary protein excretion and renal function [134]. Histological examination of the kidney showed that animals given the thromboxane synthase inhibitor were also significantly protected from glomerulosclerosis. These findings are consistent with those of others [110, 126] in other experimental models of progressive nephropathy (that is, Milan normotensive rats and NZB/NZW F₁ mice) in which a protec-

tive effect of thromboxane synthase inhibitors on renal injury was evident.

Manipulating the thromboxane biosynthetic pathway in renal disease may also be beneficial because it limits platelet activation, one of the possible mechanisms involved in the progression of renal disease. This possibility is consistent with an observational study that suggested that antiplatelet agents slowed the deterioration of renal function and prevented the development of end-stage renal disease in patients with membranoproliferative glomerulonephritis [135]. Intraglomerular platelet aggregation could facilitate mesangial cell proliferation and excessive generation of mesangial matrix. Thus, platelet-derived growth factor (PDGF) binds to specific receptors on cultured mesangial cells and stimulates DNA synthesis and cellular proliferation [136]. In vitro studies have documented that another platelet product, transforming growth factor- β (TGF- β) stimulates synthesis of the extracellular matrix components, collagen IV and laminin and the proteoglycans, biglycan and decorin, by cultured mesangial cells [137, 138] without stimulation of cell proliferation. TGF- β mediates proteoglycan production and mesangial matrix expansion in vivo in a rat model of glomerulonephritis induced by anti-thymocyte serum [139].

An alternative possibility to explain why reducing renal TxA₂ synthesis is protective against glomerulosclerosis emerged from studies showing that PGH₂/TxA₂ analogs induce a dose-dependent increase in laminin A, B₁ and B₂ chains in tumor cells in culture and that these effects are reversed by a thromboxane receptor antagonist [140]. This possibility is further supported by the recent observation that a thromboxane synthase inhibitor ameliorated progressive diabetic nephropathy and prevented the increase in type IV collagen mRNA levels in mice with non-insulin dependent diabetes mellitus [141].

Implications of pharmacologic studies

PGH₂ and TxA₂ exert multiple actions within the kidney which may contribute to the progression of renal disease. These include constriction of glomerular afferent and efferent arterioles, intraglomerular platelet aggregation, contraction of glomerular mesangial cells and proliferation of mesangial cell matrix. These effects have been characterized using stable PGH₂/TxA₂ analogs and although radioligand studies suggest a single class of specific binding sites in glomeruli, it is unclear whether these are subtypes distinct from those in platelets and vascular smooth muscle cells and whether their affinity for PGH₂ and TxA₂ is similar. In this regard, much of the information suggesting the functional importance of this system has been obtained with specific inhibitors of thromboxane synthase, an intervention that might be expected to lead to accumulation of PGH₂, coincidental with inhibition of TxA₂ synthesis.

Such studies have implicated TxA₂ in the nephrotoxicity of pharmacologic, immune and toxic insults, in the progression of chronic renal disease and in renal allograft rejection. Biochemical assessment of renal biosynthesis of TxA₂ has facilitated the identification of potential targets for pharmacological intervention. This evidence has been accumulated particularly in animal models, but also, to a limited extent, in humans. It suggests that pharmacological inhibition of TxA₂ synthesis or action merits

Table 2. Thromboxane A₂ in experimental and human renal diseases

Disease model	Site of abnormal TxA ₂ synthesis	Changes in urinary excretion of TxA ₂ metabolites	Effect of pharmacological inhibition of TxA ₂ synthase/receptor
Experimental			
Endotoxin-induced ARF (rat)	cortex [118]		↓ proteinuria, ↑ GFR and RPF [74]
Nephrotoxic nephritis (rat)	glomeruli [74]		↑ GFR [80]
Glycerol-induced ARF (rat)	glomeruli [79]		↓ proteinuria [102]
Adriamycin nephrosis (rat)	glomeruli [102]	↑ TxB ₂ [102]	↓ proteinuria [103]
Puromycin nephrosis (rat)			↓ proteinuria [105]
Passive Heymann nephritis (rat)			
STZ-induced diabetes (rat)		↑ TxB ₂ [107]	↓ albuminuria, ↑ GFR [107]
Milan normotensive strain (rat)	glomeruli [119]		↓ proteinuria [110]
Nephrotoxic nephritis (rabbit)	glomeruli [112]		↓ proteinuria, ↑ GFR [112]
CsA nephrotoxicity (rat)	glomeruli [121]	↑ TxB ₂ [119, 120, 121] ↑ 2,3-dinor-TxB ₂ [120, 121]	↑ GFR [119]
Endotoxin-induced ARF (rat)	cortex [122]		↑ GFR and RPF [122]
Ureteral obstruction (rat, rabbit)	kidney microsomes [123]		↑ GFR and RBF [124]
Renal allograft rejection (rat)			↑ GFR and RPF [128]
Renal mass ablation (rat)		↑ TxB ₂ [131]	↓ proteinuria, ↑ GFR and RPF, ↓ FSG [131]
Dahl-S strain (rat)	glomeruli [134]		↓ proteinuria, ↑ GFR, ↓ FSG [134]
Noninsulin-dependent diabetes (mouse)			↓ cortex type IV collagen mRNA [141]
Lupus nephritis (mouse)	cortex [126, 127] medulla [126]		↓ GFR [127]
Clinical			
Lupus nephritis		↑ TxB ₂ [88]	↑ GFR and RPF [88]
Nephrotic syndrome		↑ 2,3-dinor-TxB ₂ [106]	↓ proteinuria [106]
CsA nephrotoxicity		↑ 2,3-dinor-TxB ₂ [118]	

continued evaluation in the treatment of a variety of renal diseases in humans.

Because PGH₂ shares with TxA₂ many of the renal actions that have been discussed in this review, it is puzzling that pharmacological blockade of thromboxane synthase, which results in accumulation of PGH₂ in platelets [142, 143], was shown so consistently effective in ameliorating TxA₂-dependent loss of renal function and structure. This is at variance with the rather disappointing results obtained with thromboxane-synthase inhibitors when used as potential antithrombotic agents [144]. A number of studies both in vitro [145] and in vivo [146, 147] have demonstrated that PG endoperoxides modulate the functional response of platelets to thromboxane synthase inhibition and may, in fact, substitute for the proaggregatory activity of TxA₂.

Several factors might contribute to this apparent discrepancy. These include: (a) a different biosynthetic capacity of platelets, vis-à-vis glomerular cells, in producing TxA₂, resulting in different levels of "critical" pharmacological suppression; (b) a lower affinity of PGH₂ than TxA₂ for the putatively shared glomerular mesangial and arteriolar receptor(s); (c) a markedly different metabolic fate of PGH₂ in platelets (largely converted to TxA₂ by thromboxane synthase) versus glomerular mesangial and epithelial cells (perhaps largely converted to PGE₂ and PGI₂ by other isomerases) resulting in much greater accumulation of PGH₂ in platelets than in glomerular cells, in the face of comparable blockade of thromboxane synthase. Whether PGH₂/TxA₂-receptor antagonists may mimic or enhance the renal effects of thromboxane synthase inhibitors

largely remains to be investigated and may shed some light on this controversial issue.

Summary and conclusions

PGH₂ and TxA₂ exert their actions via tissue specific, receptor isoforms. PGH₂/TxA₂-dependent platelet aggregation and contraction of vascular and bronchial smooth muscle and of glomerular mesangial cells occur via receptors linked to activation of phospholipase C.

Although PGH₂/TxA₂ appear to be of little importance in the maintenance of renal function under physiological circumstances, increased renal TxA₂ biosynthesis has been documented in a variety of animal models of renal disease and in some clinical disorders (Table 2). The effects of this eicosanoid on renal tissues in vitro and of pharmacological manipulation of TxA₂ synthesis and action in vivo suggest that such interventions will provide new drugs for the treatment of human kidney disease.

GIUSEPPE REMUZZI,
GARRET A. FITZGERALD, and
CARLO PATRONO
Bergamo, Italy,
Dublin, Ireland, United Kingdom and
Chieti, Italy

Acknowledgments

This review was supported by grants from the National Institutes of Health (HL30400 and GM15431) and the Wellcore Trust. We are indebted to E. Stuart, M.L. Bonanomi and L. Piccoli for editorial

assistance. The helpful cooperation of Drs. N. Perico, C. Zoja and A. Benigni is also gratefully acknowledged.

Reprint requests to Garret A. FitzGerald, M.D., Center for Cardiovascular Science, Department of Medicine and Experimental Therapeutics, University College Dublin, 41 Eccles St., Dublin 7, Ireland.

References

- HAMBERG M, SVENSSON J, SAMUELSSON B: Thromboxanes: A new group of biologically active compounds derived from prostaglandin endoperoxides. *Proc Natl Acad Sci USA* 72:2994-2998, 1975
- ELLIS EF, OELZ O, ROBERTS LJ II, PAYNE NA, SWEETMAN BJ, NIES AS, OATES JA: Coronary arterial smooth muscle contraction by a substance released from platelets: Evidence that it is thromboxane A_2 . *Science* 193:1135-1137, 1976
- MENÉ P, DUNN MJ: Contractile effects of TxA_2 and endoperoxide analogues on cultured rat glomerular mesangial cells. *Am J Physiol* 251:F1029-1035, 1986
- MURRAY R, FITZGERALD GA: Regulation of thromboxane receptor activation in human platelets. *Proc Natl Acad Sci USA* 86:124-128, 1989
- DORN GW, II, SENS D, CHAIKHOUNI A, MAIS D, HALUSHKA PV: Cultured human vascular smooth muscle cells with functional thromboxane A_2 receptors: Measurements of U46619-induced 45 calcium efflux. *Cir Res* 60:952-956, 1987
- MENÉ P, DUBYAK GR, ABOUD HE, SCARPA A, DUNN MJ: Phospholipase C activation by prostaglandins and thromboxane A_2 in cultured mesangial cells. *Am J Physiol* 255:F1058-F1069, 1988
- NATHAN CF: Secretory products of macrophages. *J Clin Invest* 79:319-326, 1987
- SCHLONDORFF D: Renal prostaglandin synthesis: Sites of production and specific actions of prostaglandins. *Am J Med* 81(2B):1-11, 1986
- HENNEKENS CH, BURING JE, SANDERCOCK P, COLLINS R, PETO R: Aspirin and other antiplatelet agents in the secondary and primary prevention of cardiovascular disease. *Circulation* 80:749-756, 1989
- ROTH GJ, STANFORD N, MAJERUS PW: Acetylation of prostaglandin synthase by aspirin. *Proc Natl Acad Sci USA* 72:3073-3076, 1975
- DE WITT DL, SMITH WL: Primary structure of prostaglandin G/H synthase from sheep vesicular gland determined from the complementary DNA sequence. *Proc Natl Acad Sci USA* 85:1412-1416, 1988
- LEWIS HD, DAVIS JW, ARCHIBALD DG, PHIL M, STEINKE WE, SMITHERMAN TC, DOHERTY JE, SCHNAPER HW, LE WINTER MM, LINARES E, POUGET JM, SABHARWAL SC, CHESLER E, DE MOTIS H: Protective effects of aspirin against acute myocardial infarction and death in men with unstable angina. *N Engl J Med* 309:396-401, 1983
- CAIRNS JA, GENT M, SINGER J, FINNIE KJ, FROGATT GM, HOLDER DA, JABLONSKY G, KOSTUK WJ, McLENDEZ LJ, MYERS MG, SACKETT DL, SEALY BJ, TANSWER PH: Aspirin, sulfinpyrazone or both in unstable angina: Results of a Canadian multicenter trial. *N Engl J Med* 313:1369-1375, 1985
- THEROUX P, QUIMET H, McCANS J, LATOUR JG, JOLY P, LEVY G, PELLETIER E, JUNEAU M, STASIAK J, DE GUISE P, PELLETIER GB, RINZLER D, WATERS DD: Aspirin, heparin or both to treat acute unstable angina. *N Engl J Med* 319:1105-1111, 1988
- ISIS-2 COLLABORATIVE GROUP: Randomized trial of intravenous streptokinase, oral aspirin, both or neither among 17,187 cases of suspected acute myocardial infarction: ISIS-2. *Lancet* ii:349-360, 1988
- FITZGERALD DJ, ROY L, CATELLA F, FITZGERALD GA: Platelet activation in unstable coronary disease. *N Engl J Med* 315:983-989, 1986
- BHAGWAT SS, HAMANN PR, STILL WC, BUNTING S, FITZPATRICK FA: Synthesis and structure of the platelet aggregation factor thromboxane A_2 . *Nature* 315:511-513, 1985
- JOHN RA: Synthesis of thromboxane, prostacyclin and the endoperoxides, in *Advances in Prostaglandin, Thromboxane and Leukotriene Research* (vol 14) edited by PIKE JE, MORTON DR, New York, Raven Press, 1985, pp 131-154
- LEFER AM, SMITH EF III, ARAKI H, SMITH JB, AHARONY D, CLAREMON DA, MAGOLDA RL, NICOLAOU KC: Dissociation of vasoconstrictor and platelet aggregatory activities of thromboxane by carbocyclic thromboxane A_2 , a stable analog of thromboxane A_2 . *Proc Natl Acad Sci USA* 77:1706-1710, 1980
- MORINELLI TA, OKWU AK, MAIS DE, HALUSHKA PV, JOHN V, CHEN C-K, FRIED J: Difluorothromboxane A_2 and stereoisomers: Stable derivatives of thromboxane A_2 with differential effects on platelets and blood vessels. *Proc Natl Acad Sci USA* 86:5600-5604, 1989
- MAYEUX PR, MORTON HE, GILLARD J, LORD A, MORINELLI TA, BOEHM A, MAIS DE, HALUSHKA PV: The affinities of prostaglandin H_2 and thromboxane A_2 for their receptor are similar in washed human platelets. *Biochem Biophys Res Commun* 157:733-739, 1988
- LUMLEY P, WHITE BP, HUMPHREY PPA: GR32191, a highly potent and specific thromboxane A_2 receptor blocking drug on platelets and vascular and airways smooth muscle *in vitro*. *Br J Pharmacol* 97:783-794, 1989
- TAKAHARA K, MURRAY R, FITZGERALD GA, FITZGERALD DJ: The response to thromboxane A_2 in human platelets: Discrimination of two binding sites linked to distinct effector systems. *J Biol Chem* 265:6836-6844, 1990
- DORN GW II: Distinct platelet thromboxane A_2 /prostaglandin H_2 receptor subtypes. A radioligand binding study of human platelets. *J Clin Invest* 84:1883-1891, 1989
- PATSCHKE H: Pseudo-irreversible antagonism produced by the TxA_2 /PGH $_2$ receptor antagonist daltroban. (abstract) 7th International Conference on Prostaglandins and Related Compounds, Florence, May 29-June 1, 1990; p 33
- CARMO LG, HATMI M, ROTILIO D, VARGAFTIG BB: Platelet desensitization induced by arachidonic acid is not due to cyclooxygenase inactivation and involves the endoperoxide receptor. *Br J Pharmacol* 85:849-859, 1985
- HIRATA M, HAYASHI Y, USHUKUBI F, YOKOTA Y, KAGEYAMA R, NAKANISHI S, NARUMIYA S: Cloning and expression of cDNA for a human thromboxane A_2 receptor. *Nature* 349:617-620, 1991
- LEDUC L, WYCHE A, SPRECHER H, SANKARAPPE S, NEEDLEMAN P: Analogues of arachidonic acid used to evaluate structural determinants of prostaglandin receptor and enzyme specificities. *Mol Pharmacol* 19:242-247, 1981
- MAIS DE, DEHOLL D, SIGHTLER H, HALUSHKA PV: Different pharmacologic activities for 13-azapinane thromboxane A_2 analogs in platelets and blood vessels. *Eur J Pharm* 148:309-315, 1988
- MAIS DE, SAUSSY DL, CHAIKHOUNI A, KOCHER PJ, KNAPP DR, HAMANAKI N, HALUSHKA PV: Pharmacologic characterization of human and canine thromboxane A_2 /prostaglandin H_2 receptors in platelets and blood vessels: Evidence for different receptors. *J Pharm Exp Ther* 233:418-424, 1985
- JONES RL, PEESAPATA V, WILSON NH: Antagonism of the thromboxane-sensitive contractile systems of the rabbit aorta, dog saphenous vein and guinea-pig trachea. *Br J Pharmacol* 76:423-429, 1982
- AKBAR H, MUKHOPADHYAY A, ANDERSON K, NAVRAN S, RAMSTEDT K, MILLER D, FELLER D: Antagonism of prostaglandin-mediated responses in platelets and vascular smooth muscle by 13-azaprostanoic acid analogs. *Biochem Pharmacol* 34:641-647, 1985
- SWAYNE GTG, MAGUIRE J, DOLAN J, RAVAL R, DANE G, GREENER M, OWEN DAA: Evidence for homogeneity of thromboxane A_2 receptors using structurally different antagonists. *Eur J Pharmacol* 152:311-319, 1988
- FURCI L, MURRAY R, FITZGERALD DJ, FITZGERALD GA: Heterogeneity of prostaglandin H_2 /thromboxane A_2 receptors: Distinct subtypes mediate vascular smooth muscle contraction and platelet aggregation. *J Pharm Exp Ther* 258:74-81, 1991
- MURRAY R, SHIPP L, FITZGERALD GA: PGH $_2$ /thromboxane receptor desensitization: Cross talk with adenylate cyclase in human platelets. *J Biol Chem* 265:21670-21675, 1990

36. MENÈ P, ABBOUD HE, DUNN MJ: Regulation of human mesangial cell growth in culture by thromboxane A₂ and prostacyclin. *Kidney Int* 38:232-239, 1990
37. BRASS LF, SHALLER CC, BELMONTE EJ: Inositol 1,4,5-triphosphate-induced granule secretion in platelets. Evidence that the activation of phospholipase C mediated by platelet thromboxane receptors involves a guanine nucleotide binding protein-dependent mechanism distinct from that of thrombin. *J Clin Invest* 79:1269-1275, 1987
38. BRASS L, LAPOSATA M, BANGA H, RITTENHOUSE S: Regulation of the phosphoinositide hydrolysis pathway in thrombin-stimulated platelets by a pertussis toxin-sensitive guanine nucleotide binding protein. *J Biol Chem* 261:16838-16847, 1986
39. GUPTA SK, DIEZ E, HEASLEY LE, OSAWA S, JOHNSON GL: A G protein mutant that inhibits thrombin and purinergic receptor activation of phospholipase A₂. *Science* 662-666, 1990
40. HOUSLAY M, BOJANIC D, WILSON A: Platelet activating factor and U46619 stimulate a GTPase activity in human platelets which is distinct from the guanine nucleotide regulatory proteins. *Biochem J* 234:737-740, 1986
41. JEREMY JY, MIKHAILIDIS DP, DANDONA P: Thromboxane A₂ analogue (U-46619) stimulates vascular PGI₂ synthesis. *Eur J Pharm* 107:259-262, 1985
42. NARUMIYA S, USHIKUBI F, NAKAJIMA N, HIRATA M, OKUMA M: Isolation and characterization of the thromboxane A₂/prostaglandin H₂ receptor from human blood platelets. (abstract) 7th International Conference on Prostaglandins and Related Compounds 1990; Florence, May 29-June 1, p. 199
43. WILKES B, MENTO P, MACICA C, SOLOMON J: Reduced thromboxane receptors in diabetic glomeruli: Relationship to hyperfiltration. (abstract) *Kidney Int* 35:336, 1989
44. SCHAMBELAN M, BLAKE S, SRAER J, BENS M, NIVEZ MP, WAHBE F: Increased prostaglandin production by glomeruli isolated from rats with streptozotocin-induced diabetes mellitus. *J Clin Invest* 75:404-412, 1985
45. SMITH WL, BELL TG: Immunohistochemical localization of the prostaglandin-forming cyclo-oxygenase in renal cortex. *Am J Physiol* 235:451-457, 1978
46. FOLKERT VW, SCHLONDORFF D: Prostaglandin synthesis in isolated glomeruli. *Prostaglandins* 17:79-86, 1979
47. HASSID A, KONIECZKOWSKY M, DUNN MJ: Prostaglandin synthesis in isolated rat glomeruli. *Proc Natl Acad Sci USA* 76:1155-1159, 1979
48. SRAER J, ARDAILLOU N, SRAER JD, ARDAILLOU R: *In vitro* prostaglandin synthesis by human glomeruli and papillae. *Prostaglandins* 23:855-864, 1982
49. STAHL RA, PARAVICINI M, SCOLLMEYER P: Angiotensin II stimulation of prostaglandin E₂ and 6-keto PGF_{1α} formation by isolated human glomeruli. *Kidney Int* 26:30-34, 1984
50. LIVIO M, CHIABRANDO C, MACCONI D, BENIGNI A, ZIMEI M, DEPIETRO MT, REMUZZI G: Metabolism of arachidonic acid in isolated glomeruli from pig kidney. *Biochem Biophys Acta* 961:110-121, 1988
51. PETRULIS AS, AIKAWA M, DUNN MJ: Prostaglandin and thromboxane synthesis by rat glomerular epithelial cells. *Kidney Int* 20:469-474, 1981
52. SCHARSCHMIDT LA, DUNN MJ: Prostaglandin synthesis by rat glomerular mesangial cells in culture. Effects of angiotensin II and arginine vasopressin. *J Clin Invest* 71:1756-1764, 1983
53. FOLKERT VW, YUNIS M, SCHLONDORFF D: Prostaglandin synthesis linked to phosphatidylinositol turnover in isolated rat glomeruli. *Biochem Biophys Acta* 794:206-217, 1984
54. SIMONSON MS, DUNN MJ: Endothelin-1 stimulates contraction of rat glomerular mesangial cells and potentiates β-adrenergic-mediated cyclic adenosine monophosphate accumulation. *J Clin Invest* 85:790-797, 1990
55. ZOJA C, BENIGNI A, RENZI D, PICCINELLI A, PERICO N, REMUZZI G: Endothelin and eicosanoid synthesis in cultured mesangial cells. *Kidney Int* 37:927-933, 1990
56. SCHUMACHER WA, STEINBACHER TE, ALLEN GT, OGLETREE ML: Role of thromboxane receptor activation in the bronchospastic response to endothelin. *Prostaglandins* 40:71-79, 1990
57. REYNOLDS EE, MOK LLS: Role of thromboxane A₂/prostaglandin H₂ receptor in the vasoconstrictor response of rat aorta to endothelin. *J Pharm Exp Ther* 252:915-921, 1990
58. STORK JE, SHEN TY, DUNN MJ: Stimulation of prostaglandin E₂ and thromboxane B₂ production in cultured rat mesangial cells by platelet activating factor: Inhibition by a specific receptor antagonist. (abstract) *Kidney Int* 27:267, 1985
59. LOVETT DH, RESCH K, GEMSA D: Interleukin 1 and the glomerular mesangium. II. Monokine stimulation of mesangial cell prostanoic secretion. *Am J Pathol* 129:543-551, 1987
60. RAZ A, WYCHE A, NEEDLEMAN P: Temporal and pharmacological division of fibroblast cyclooxygenase expression into transcriptional and translational phases. *Proc Natl Acad Sci USA* 86:1657-1661, 1989
61. MALER JAM, HLA T, MACIAG T: Cyclooxygenase is an immediate-early gene induced by interleukin-1 in human endothelial cells. *J Biol Chem* 265:10805-10808, 1990
62. BAUD L, NIVEZ M-P, CHANSEL D, ARDAILLOU R: Stimulation by oxygen radicals of prostaglandin production by rat renal glomeruli. *Kidney Int* 20:322-339, 1981
63. ADLER S, STAHL AK, BAKER PJ, CHEN YP, PRITZ PM, COUSER WG: Biphasic effect of oxygen radicals on prostaglandin production by rat mesangial cells. *Am J Physiol* 252:F743-F749, 1987
64. HANSCH GM, BETZ M, GUNTHER J, ROTHER KO, STERZEL B: The complement membrane attack complex stimulates the prostanoic production of cultured glomerular epithelial cells. *Int Arch Allergy Appl Immunol* 85:87-93, 1988
65. WANG J, DUNN MJ: Endotoxin stimulates prostaglandin release by cultured rat glomerular mesangial cells. (abstract) *Kidney Int* 31:291, 1987
66. KNAUSS T, MENÈ P, RICANATI SA, KESTER M, DUBYAK GR, EMANCIPATOR SN, SEDOR JR: Immune complex activation of rat glomerular mesangial cells: Dependence on the Fc region of antibody. *Am J Physiol* 257:F478-F485, 1989
67. CHEN Y-P, STAHL RAK, ADLER S, BAKER P, PRITZL P, COUSER WG: Antibody (Ab) to a mesangial cell (MC) membrane antigen stimulates prostaglandin production by a complement (C)-dependent mechanism. (abstract) *Kidney Int* 31:315, 1987
68. SRAER J, WOLF C, OUDINET J-P, BENS M, ARDAILLOU R, SRAER J-D: Human glomeruli release fatty acids which stimulate thromboxane synthesis in platelets. *Kidney Int* 32:62-68, 1987
69. BAYLIS C: Effects of administered thromboxane on the intact, normal rat kidney. *Renal Physiol* 10:110-121, 1987
70. FEIGEN LP, CHAPNICK BM, FLEMMING JE, FLEMMING JM, KADOWITZ PJ: Renal vascular effects of endoperoxide analogs, prostaglandins and arachidonic acid. *Am J Physiol* 233:H573-H579, 1977
71. GERBER JG, ELLIS E, HOLLIFIELD J, NIES AS: Effect of prostaglandin endoperoxide analogue on canine renal function. *Eur J Pharmacol* 53:239-246, 1979
72. LOUTZENHISER R, EPSTEIN M, HORTON C, SONKE P: Reversal of renal and smooth muscle actions on the thromboxane mimetic U-44069 by diltiazem. *Am J Physiol* 250:F619-F626, 1986
73. EPSTEIN M, HAYASHI K, LOUTZENHISER R: Direct evidence that thromboxane mimetic U44069 preferentially constricts the afferent arteriole. (abstract) *Kidney Int* 35:291, 1989
74. LIANOS EA, ANDRES GA, DUNN MJ: Glomerular prostaglandin and thromboxane synthesis in rat nephrotoxic nephritis. Effects on renal hemodynamics. *J Clin Invest* 72:1439-1448, 1986
75. BLANTZ RC, WILSON CB: Acute effects of antiglomerular basement membrane antibody on the process of glomerular filtration in the rat. *J Clin Invest* 58:899-911, 1976
76. SCHARSCHMIDT LA, LIANOS E, DUNN MJ: Arachidonate metabolites and the control of glomerular function. *Fed Proc* 42:3058-3063, 1983
77. CHATZANTONIOU C, DANIELS FH, ARENDSHORST WJ: Exaggerated renal vascular reactivity to angiotensin and thromboxane in young genetically hypertensive rats. *Am J Physiol* 259:F372-F382, 1990
78. ARDAILLOU N, HAGEGE J, NIVEZ M-P, ARDAILLOU R, SCHLONDORFF D: Vasoconstrictor-evoked prostaglandin synthesis in cultured human mesangial cells. *Am J Physiol* 248:F240-F246, 1985
79. SRAER J-D, MOULONGUET-DOLERIS L, DELARUE F, SRAER J, ARDAILLOU R: Prostaglandin synthesis by glomeruli isolated from

- rats with glycerol-induced acute renal failure. *Circ Res* 49:775-783, 1981
80. PAPANICOLAOU N, HATZIANTONIOU C, BARIETY J: Selective inhibition of thromboxane synthesis partially protected while inhibition of angiotensin II formation did not protect rats against acute renal failure induced with glycerol. *Prostaglandin Leuk Med* 21:29-35, 1986
 81. BLASINGHAM MC, NASJILETTI A: Differential effects of cyclooxygenase inhibition in sodium replete and sodium-deprived dog. *Am J Physiol* 239:F360-F365, 1980
 82. BLASINGHAM MC, SHADE RE, SHARE L, NASJILETTI A: The effect on meclofenamate on renal blood flow in the unanesthetized dog: Relation to renal prostaglandins and sodium balance. *J Pharm Exp Ther* 214:1-4, 1980
 83. DE FORREST JM, DAVIS JO, FREEMAN RH, SEYMOUR AA, ROWE BP, WILLIAMS GM, DAVIS TP: Effects of indomethacin and meclofenamate on renin release and renal hemodynamic function during chronic sodium depletion in conscious dogs. *Circ Res* 47:99-107, 1980
 84. KIRSCHENBAUM MA, WHITE N, STEIN JH, FERRIS TF: Redistribution of renal cortical blood flow during inhibition of prostaglandin synthesis. *Am J Physiol* 227:801-805, 1974
 85. HEINRICH WL, ANDERSON RL, BERNIS AL: The role of renal nerves and prostaglandins in control of renal hemodynamics and plasma renin activity during hypotensive hemorrhage in the dog. *J Clin Invest* 61:744-750, 1978
 86. OLIVER JA, SCIACCA RR, PINTO J, CANNON PJ: Participation of the prostaglandins in the control of renal blood flow during acute reduction of cardiac output in the dog. *J Clin Invest* 67:229-237, 1981
 87. ZIPSER RD: Effects of selective inhibition of thromboxane synthesis on renal function in humans. *Am J Physiol* 248:F753-F756, 1985
 88. PIERUCCI A, SIMONETTI BM, PECCI G, MAVRIKAKIS G, FERIOZZI S, CINOTTI GA, PATRIGNANI P, CIABATTONI G, PATRONO C: Improvement of renal function with selective thromboxane antagonism in lupus nephritis. *N Engl J Med* 320:421-425, 1989
 89. FITZGERALD GA, PEDERSEN AK, PATRONO C: Analysis of prostacyclin and thromboxane biosynthesis in cardiovascular disease. *Circulation* 67:1174-1177, 1983
 90. PATRONO C, CIABATTONI G, PUGLIESE F, PIERUCCI A, BLAIR IA, FITZGERALD GA: Estimated rate of thromboxane secretion into the circulation of normal man. *J Clin Invest* 77:590-594, 1986
 91. ROBERTS LJ, II, SWEETMAN BJ, OATES JA: Metabolism of thromboxane B₂ in man. Identification of twenty urinary metabolites. *J Biol Chem* 256:8384-8393, 1981
 92. FITZGERALD GA, BRASH AR, BLAIR IA, LAWSON J: Analysis of urinary metabolites of thromboxane and prostacyclin by negative ion-chemical ionization gas chromatography-mass spectrometry, in *Advances in Prostaglandin Thromboxane and Leukotriene Research 15*, edited by HAYASHI O, YAMAMOTO S, New York, Raven Press, 1985, pp. 87-90
 93. PATRONO C: Measurement of thromboxane biosynthesis in man. *Eicosanoids* 2:249-251, 1989
 94. CATELLA F, HEALY D, LAWSON J, FITZGERALD GA: 11-dehydrothromboxane B₂: An index of thromboxane formation in the human circulation. *Proc Natl Acad Sci USA* 83:5861-5865, 1986
 95. CIABATTONI G, BOSS AH, PATRIGNANI P, CATELLA F, SIMONETTI BM, PIERUCCI A, PUGLIESE F, FILABOZZI P, PATRONO C: Effects of sulindac on renal and extrarenal eicosanoid synthesis. *Clin Pharmacol Ther* 41:380-383, 1987
 96. FITZGERALD GA, OATES JA, HAWIGER J, MAAS RL, ROBERTS LJ, BRASH AR: Endogenous synthesis of prostacyclin and thromboxane and platelet function during chronic aspirin administration in man. *J Clin Invest* 71:676-688, 1983
 97. PATRONO C, CIABATTONI G, REMUZZI G, GOTTI E, BOMBARDIERI S, DiMUNNO O, TARTARELLI G, CINOTTI GA, SIMONETTI BM, PIERUCCI A: Functional significance of renal prostacyclin and thromboxane A₂ production in patients with systemic lupus erythematosus. *J Clin Invest* 76:1011-1015, 1985
 98. BENIGNI A, CHIABRANDO C, PERICO N, FANELLI R, PATRONO C, FITZGERALD GA, REMUZZI G: Renal metabolism and urinary excretion of thromboxane B₂ in the rat. *Am J Physiol* 257:F77-F85, 1989
 99. REILLY IAG, DORAN J, SMITH B, FITZGERALD GA: Increased thromboxane biosynthesis in a human model of platelet activation: Biochemical and functional consequences of selective inhibition of thromboxane synthase. *Circulation* 73:1300-1309, 1986
 100. O'SHAUGHNESSY K, WARD P, TAYLOR I, TAYLOR G, FULLER R: Urinary excretion of thromboxane B₂ and its dinor metabolite in normal subjects following inhalation and ingestion of thromboxane B₂. (abstract) *Am Rev of Resp Dis* 141(4):A392, 1990
 101. PATRONO C, SALVATI P: Thromboxane biosynthesis and pharmacologic modulation in progressive glomerulosclerosis. *Am J Nephrol* 9 (Suppl 1):13-16, 1989
 102. REMUZZI G, IMBERTI L, ROSSINI M, MORELLI C, CARMINATI C, CATTANEO GM, BERTANI T: Increased glomerular thromboxane synthesis as a possible cause of proteinuria in experimental nephrosis. *J Clin Invest* 75:94-101, 1985
 103. GOTO T, MUNE M, MATOBA K, YUKAWA S, NAMOTO H: Effects of selective thromboxane A₂ synthetase inhibitor on aminonucleoside induced nephrotic rats. (abstract) *Xth International Congress of Nephrology*, London, July 26-31, 1987, p 227
 104. SUZUKI J, SUZUKI S, SATO K, WATANABE H, KAWASAKI Y, NOZAWA R, KUMAKI S, KUME K, HIGUCHI E, KAMIYAMA S, YUGETA E, KATO K, SUZUKI H: The role of thromboxane A₂ in aminonucleoside nephrotic rats. (abstract) *8th Congress of the International Pediatric Nephrology Association*, Toronto, 1989
 105. CYBULSKY AV, LIEBERTHAL W, QUIGG RJ, RENNKE HG, SALANT DJ: A role for thromboxane in complement-mediated glomerular injury. *Am J Pathol* 128:45-51, 1987
 106. NIWA T, MAEDA K, SHIBATA M, YAMADA K: Clinical effects of selective thromboxane A₂ synthetase inhibitor in patients with nephrotic syndrome. *Clin Nephrol* 30:276-281, 1988
 107. CRAVEN PA, DERUBERTIS FR: Thromboxane synthetase inhibition suppresses urinary albumin excretion (UAlb) in diabetic rats. (abstract) *Kidney Int* 37:347, 1990
 108. COLLINS DM, MANISCALCO M, BEST CF, COFFMAN TM, KLOTMAN E: A thromboxane synthetase inhibitor (TSI) ameliorates the proteinuria induced by high protein feeding in diabetic rats. (abstract) *Kidney Int* 37:501, 1990
 109. BRANDIS A, BIANCHI G, REALE E, HELMCHEN U, KUHN K: Age-dependent glomerulosclerosis and proteinuria occurring in rats of the Milan normotensive strain and not in rats of the Milan hypertensive strain. *Lab Invest* 55:234-243, 1986
 110. SALVATI P, FERTI C, FERRARIO RG, LAMBERTI E, DUZZI L, BIANCHI G, REMUZZI G, PERICO N, BENIGNI A, BRAIDOTTI P, COGGI G, PUGLIESE F, PATRONO C: Role of enhanced glomerular synthesis of thromboxane A₂ in progressive kidney disease. *Kidney Int* 38:447-458, 1990
 111. LIANOS EA, ANDRES GA, DUNN MJ: Glomerular prostaglandin and thromboxane synthesis in rat nephrotic serum nephritis. Effects on renal hemodynamics. *J Clin Invest* 72:1439-1448, 1983
 112. MACCONI D, BENIGNI A, MORIGI M, UBIALI A, ORISIO S, LIVIO M, PERICO N, BERTANI T, REMUZZI G, PATRONO C: Enhanced glomerular thromboxane A₂ mediates some pathophysiologic effect of platelet-activating factor in rabbit nephrotoxic nephritis: Evidence from biochemical measurements and inhibitor trials. *J Lab Clin Med* 113:549-560, 1989
 113. REMUZZI A, BATTAGLIA C, ROSSI L, ZOJA C, REMUZZI G: Glomerular size-selectivity in nephrotic rats exposed to diets with different protein content. *Am J Physiol* 253:F318-F327, 1987
 114. REMUZZI A, BATTAGLIA C, REMUZZI G: Effect of low-protein diet on size and charge permeability in experimental nephrosis. (abstract) *Kidney Int* 37:472, 1989
 115. FOGO A, YOSHIDA Y, GLICK AD, HOMMA T, ICHIKAWA I: Serial micropuncture analysis of glomerular function in two rat models of glomerular sclerosis. *J Clin Invest* 82:322-330, 1988
 116. SCHOLEY JW, MILLER PL, RENNKE HG, MEYER TW: Effect of converting enzyme inhibition on the course of adriamycin-induced nephropathy. *Kidney Int* 36:816-822, 1989
 117. YOO J, SCHLONDORFF D, NEUGARTEN J: Thromboxane mediates the renal hemodynamic effects of platelet activating factor. *J Pharmacol Exp Ther* 253:743-748, 1990
 118. KUHN K, FORSTERMANN U, FROLICH JC, VETERQVIST O, GREEN

- K, BRUNKHORST R, KOCH KM, HAAS J: Effect of cyclosporin A (CyA) on blood pressure and prostacyclin and thromboxane A₂ production, in *Abstracts, Xth International Congress of Nephrology*, London 26-31, July 1987, p 231
119. PERICO N, BENIGNI A, ZOJA C, DELAINI F, REMUZZI G: Functional significance of exaggerated renal thromboxane A₂ synthesis induced by cyclosporin A. *Am J Physiol* 251:F581-F587, 1986
120. COFFMAN TM, CARR DR, YARGER WE, KLOTMAN PE: Evidence that renal prostaglandin and thromboxane production is stimulated in chronic cyclosporine nephrotoxicity. *Transplantation* 43:282-285, 1987
121. BENIGNI A, CHIABRANDO C, PICCINELLI A, PERICO N, GAVINELLI M, FURCI L, PATINO O, ABBATE M, BERTANI T, REMUZZI G: Increased urinary excretion of thromboxane B₂ and 2,3-dinor-TxB₂ in cyclosporin A nephrotoxicity. *Kidney Int* 34:164-174, 1988
122. BADR KF, KELLEY VE, RENNKE HG, BRENNER BM: Roles for thromboxane A₂ and leukotrienes in endotoxin-induced acute renal failure. *Kidney Int* 30:474-480, 1986
123. MORRISON AR, NISHIKAWA K, NEEDLEMAN P: Unmasking of thromboxane A₂ synthesis by ureteral obstruction in the rabbit kidney. *Nature* 267:259-260, 1977
124. KLOTMAN PE, SMITH SR, VOLPP BD, COFFMAN TM, YARGER WE: Thromboxane synthetase inhibitor improves function of hydronephrotic rat kidneys. *Am J Physiol* 250:F282-F287, 1986
125. ALBRIGHTSON CR, EVERS AS, GRIFFIN AC, NEEDLEMAN P: Effect of endogenously produced leukotrienes and thromboxane on renal vascular resistance in rabbit hydronephrosis. *Circ Res* 61:514-522, 1987
126. KELLEY VE, SNEVE S, MUSINSKI S: Increased renal thromboxane production in murine lupus nephritis. *J Clin Invest* 77:252-259, 1986
127. SPURNEY RF, RUIZ P, KLOTMAN PE, PISETSKY DS, COFFMAN TM: Functional significance of thromboxane (Tx) in murine lupus nephritis (LN). (abstract) *Kidney Int* 37:351, 1990
128. COFFMAN TM, YARGER WE, KLOTMAN PE: Functional role of thromboxane production by acutely rejecting renal allografts in rats. *J Clin Invest* 75:1242-1248, 1985
129. HOSTETTER TH, OLSON JL, RENNKE HG, VENKATACHALAM MA, BRENNER BM: Hyperfiltration in remnant nephrons: A potentially adverse response to renal ablation. *Am J Physiol* 241:F85-F93, 1981
130. OLSON JJ, HOSTETTER TH, RENNKE HG, BRENNER BM, VENKATACHALAM MA: Altered glomerular permselectivity and progressive sclerosis following extreme ablation of renal mass. *Kidney Int* 22:112-116, 1982
131. PURKERSON ML, JOIST JH, YATES J, VALDES A, MORRISON A, KLAHR S: Inhibition of thromboxane synthesis ameliorates the progressive kidney disease of rats with subtotal renal ablation. *Proc Natl Acad Sci USA* 82:193-197, 1985
132. ZOJA C, BENIGNI A, LIVIO M, BERGAMELLI A, ORISIO S, ABBATE M, BERTANI T, REMUZZI G: Selective inhibition of platelet thromboxane generation with low-dose aspirin does not protect rats with reduced renal mass from the development of progressive disease. *Am J Pathol* 134:1027-1038, 1989
133. ZOJA C, PERICO N, BERGAMELLI A, PASINI M, MORIGI M, DADAN J, BELLONI A, BERTANI T, REMUZZI G: Ticlopidine prevents renal disease progression in rats with reduced renal mass. *Kidney Int* 37:934-942, 1990
134. YAMASHITA W, ITO Y, WEISS MA, OOI BS, POLLAK VE: A thromboxane synthetase antagonist ameliorates progressive renal disease of Dahl-S rats. *Kidney Int* 33:77-83, 1988
135. DONADIO JV, ANDERSON CF, MITCHELL JC, HOLLEY KE, ILSTRUP DM, FUSTER V, CHESEBRO JH: Membranoproliferative glomerulonephritis. A prospective clinical trial of platelet-inhibitor therapy. *N Engl J Med* 310:1421-1426, 1984
136. SHULTZ PJ, DiCORLETTI PE, SILVER BJ, ABBODD HE: Mesangial cells express PDGF mRNAs and proliferate in response to PDGF. *Am J Physiol* 255: (Suppl 6/part F) F674-F684, 1988
137. ISHIMURA E, STERZEL RB, KASHGARIAN M: Effect of transforming growth factor (TGF)- β on extracellular matrix (ECM) production by cultured rat mesangial cells (MCs). (abstract) *Kidney Int* 37:197, 1990
138. BORDER W, OKUDA S, LANGUINO LR, RUOSLAHTI E: Transforming growth factor- β (TGF- β) regulates production of proteoglycans by mesangial cells. *Kidney Int* 37:689-695, 1990
139. OKUDA S, LANGUINO L, RUOSLAHTI E, BORDER W: Transforming growth factor- β (TGF- β) is responsible for glomerular proteoglycan production in experimental glomerulonephritis. (abstract) *Kidney Int* 37:222, 1990
140. KLOTMAN P, BRUGGEMAN L, HASSELL J, HORIGAN E, MARTIN G, YAMADA Y: Regulation of extracellular matrix by thromboxane. (abstract) *Kidney Int* 35:294, 1989
141. LEDBETTER S, COPELAND EJ, NOONAN D, VOGELI G, HASSELL JR: Altered steady-state mRNA levels of basement proteins in diabetic mouse kidney and thromboxane synthase inhibition. *Diabetes* 39:196-203, 1990
142. MARCUS AJ, WEKSLER BB, JAFFE EA, BROEKMAN MJ: Synthesis of prostacyclin from platelet-derived endoperoxides by cultured human endothelial cells. *J Clin Invest* 66:979-986, 1980
143. NOWAK J, FITZGERALD GA: Redirection of prostaglandin endoperoxide metabolism at the platelet-vascular interface in man. *J Clin Invest* 83:380-385, 1989
144. FITZGERALD GA, REILLY IAG, PEDERSEN AK: The biochemical pharmacology of thromboxane synthase inhibition in man. *Circulation* 72:1194-1201, 1985
145. BERTELE V, FALANGA A, TOMASIAK M, CHIABRANDO C, CERLETTI C, DE GAETANO G: Pharmacologic inhibition of thromboxane synthetase and platelet aggregation: Modulatory role of cyclooxygenase products. *Blood* 63:1460-1466, 1984
146. GRESELE P, ARNOUT J, DECKMYN H, HUYBRECHTS E, PIETERS G, VERMYLEN J: Role of proaggregatory and antiaggregatory prostaglandins in hemostasis. Studies with combined thromboxane synthase inhibition and thromboxane receptor antagonism. *J Clin Invest* 80:1435-1445, 1987
147. FITZGERALD DJ, FRAGETTA J, FITZGERALD GA: Prostaglandin endoperoxides modulate the response to thromboxane synthase inhibition during coronary thrombosis. *J Clin Invest* 82:1708-1713, 1988